



Case Report

Hemodynamics, baroreflex index and blood biomarkers of a patient who died after being affected by COVID-19: case report

Valdenilson Ribeiro Ribas ^{*}, Renata de Melo Guerra Ribas ,
Mayara Guerra Souto Barros , Ketlin Helenise dos Santos Ribas ,
Nery Adamy Neto , Mayanna Guerra Souto Barros , Hugo André de Lima Martins 

Instituto do Cérebro de Pernambuco (ICerPE), Jaboatão dos Guararapes, PE, Brazil

ARTICLE INFO

Article history:

Received 12 May 2020

Accepted 15 June 2020

Available online 18 June 2020

Introduction

Baroreflex index and hemodynamics

Baroreceptors, baroreceptors, mechanoreceptors or even mechanoreceptors are receptors responsible for the contraction and dilation of blood vessels.¹ They are located mainly in the carotid sinus² and aortic arch^{3,4} and have the function of detecting variations in blood pressure and transmitting this information to the central nervous system. This information generates responses from the autonomic nervous system, modulating the functioning of the blood circulation (Baroreceptor reflex).⁵

The more the baroreceptors are activated, the more functional oxygen is transported to the lungs and the brain.⁶

This functioning with active and constant movement of these mechanoreceptors also seems to indicate that there is a good flow in the blood displacement at the level of blood microcirculation, also called hemodynamics.⁷

Adequate hemodynamics depends on the state of cardiovascular health. According to Houston (2014), the blood vessel has a finite number of responses to an infinite number of insults.⁸ Finite vascular responses are oxidative stress (Reactive Oxygen Species and Reactive Nitrogen Species),⁹ inflammation¹⁰ and exaggerated immune reaction of the organism¹¹ and are related to the pathophysiology of vascular disease, which lead to an abnormal vascular biology, such as endothelial dysfunction (ED) and vascular smooth muscle dysfunction (VSMD).⁹

In this infinite estimate of insults, there is hypertension, dyslipidemia,¹² diabetes,¹³ hyperglycemia,¹⁴ obesity,¹⁵

^{*} Corresponding author at: Instituto do Cérebro de Pernambuco (ICerPE), Galeria Piedade Center, Loja 208, av. Zequinha Barreto, 384 Piedade, Jaboatão dos Guararapes, PE CEP: 54.400-090, Brazil.

E-mail address: ribas.professor@gmail.com (V.R. Ribas).

<https://doi.org/10.1016/j.htct.2020.06.004>

2531-1379/© 2020 Associação Brasileira de Hematologia, Hemoterapia e Terapia Celular. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

smoking,¹⁶ insulin resistance,¹⁷ increased homocysteine,¹⁸ among others, which enable vascular responses with cytokines,¹⁹ chemokines,²⁰ adhesion molecules,²¹ heat shock proteins²² and antibodies,²³ leading to inflammation,²⁴ oxidative stress²⁵ and autoimmune dysfunction.²⁶

Atherosclerosis and vascular disease are postprandial phenomena, that is, they arise after eating allergenic and inflammatory foods,²⁷ associated with hyperglycemia and hypertriglyceridemia, with endothelial dysfunction being the earliest stage of vascular disease, which can be detected through the baroreflex index, hemodynamics²⁷ and biomarkers already studied in the literature, linked to cardiovascular diseases.²⁸

In this sense, this study aimed to report the values of the baroreflex and hemodynamic index related to biomarkers that signal obstruction of the blood microcirculation found in the 57-year-old male patient, who died after being affected by COVID-19.

Methods

Assessments were performed by functional neurometry (FN) and laboratory tests. The evaluation with functional neurometry was performed in an air-conditioned room at a temperature of $22 \pm 2^\circ\text{C}$. The capacity, functionality and elasticity of the blood vessels are evaluated indirectly in 3 (three) positions. Thus, the FN was performed in 3 positions: dorsal decubitus, standing up and orthostatic, called DSO analysis by the author, organizer and creator of the method, Nelson Alves Pereira Júnior.²⁷ The sympathetic autonomic nervous system represents the vasoconstrictions measured at the frequencies of 0.01–0.04 Hz and from 0.04 to 0.20 Hz and the parasympathetic represents the vasodilatations measured at the frequencies of 0.20–0.50 Hz.²⁷

The five categories of this evaluation protocol of DSO analysis were named by the author as: anxiety control; physiological response; baroreflex index; hemodynamics (blood flow velocity); and brain neurometry. Detailed descriptions of each category, which captures biological signals by sensors calibrated in Series and/or Fourier Transform algorithms, have been published in the *Journal of Psychology and Psychotherapy Research* (DOI: <https://doi.org/10.12974/2313-1047.2020.07.1>).²⁷

In this case report, the categories were approached briefly with an emphasis only on the expected values to facilitate the reader's understanding of the interpretation of the results. In Functional Neurometry, anxiety control is measured through the galvanic skin response.²⁹ Thus, the electrodermal resistance is indirectly measured by means of the sweat gland.³⁰ In this context, as sweat on the skin allows greater electrical conduction,³¹ it can be estimated that with a greater amount of sweat there seems to be less energy stored in the liver (glycogen) for a person's adrenal to enable the fight and/or flight process or for that adrenal to control anxiety through the hormones aldosterone, cortisol, adrenaline and dehydroepiandrosterone.²⁷

The anxiety control category is calculated using a scale with agreed values from 0 to 100%, in which the level of energy provided by the functional reserve already absorbed by the

enterocytes and already stored in the liver can be measured indirectly,³² awaiting the need to pass into the bloodstream by glycogenolysis³³ to make the fight and/or flight process feasible.³⁴ The minimum expected in this category is 75%.

The physiological response is measured by varying the peripheral temperature,³⁵ demonstrating whether or not there is adequate functionality of blood vessel elasticity.³⁶ This elasticity refers to an expected variation in vasoconstriction and vasodilation,³⁶ legitimizing the sympathetic (fight and flight) and parasympathetic (relaxation).³⁷

The signal origin is made through the peripheral temperature sensor, which scales the unit of measurement in degrees Celsius. The originated values generate 2 more important results, as a percentage of variability of the sympathetic and parasympathetic systems and the temperature values in the thermoregulation. Both are expressed on a scale ranging from 0 to 100 (%) or ($^\circ\text{C}$). The location of this sensor is in the proximal phalanx of the ring finger.³⁸ The expected values for an ideal thermoregulation are between 31.5°C and 32.5°C .³⁵

Baroreflex behavior is the ability of blood vessels to contract and dilate, which allows the neuroscientist to observe the level of functional oxygen that will better signal red blood cell displacement,³⁹ in the process of transporting oxygen within the blood vessels. In functional neurometry, the balance of these contractions and dilations (baroreflex index) must present a percentage above 90%.⁴⁰

Some studies have been warning about the importance of this measure of the low baroreflex index found in patients with obstructive sleep apnea,⁴¹ with slight respiratory changes⁴² asymptomatic or imperceptible, but which gradually cause mitochondrial suffering.⁴³ Other studies also demonstrate a reduction in the baroreflex index associated with anemia, inflammation,²⁷ to a lack of vitamin C,^{44,45} folic acid⁴⁶ and sedentary lifestyle⁴⁶ and may contribute to the manifestation of possible negative effects in the prevention of atherosclerosis.

Hemodynamics⁴⁷ may be directly related to endothelial dysfunction and, consequently, to the risk of cardiovascular disease.⁴⁸ In functional neurometry, the ideal hemodynamics should be below 10%. When above this percentage, it indicates a reduction in blood flow speed caused by peripheral vascular resistance, cardiac output and/or inflammatory response.²⁷

Case report

A fifty-seven-year-old male patient, assessed on February 10, 2020 at the Brain Institute of Pernambuco – *Instituto do Cérebro de Pernambuco (ICerPE)*. Before the assessment by functional neurometry, the following measurements were taken: weight [100 kg], height [1 m 78 cm], waist circumference [117 cm] and body mass index [31.6], fulfilling the classification criteria of mild obesity, with 23.3 kg overweight. The patient reported that he did not frequently do physical activity (sedentary). The patient went to ICerPE, taken by his daughter, with the objective of undergoing a complete assessment with functional neurometry and quantitative electroencephalography, after learning that there was treatment for mood disorder and generalized anxiety disorder.

When the evaluation with functional neurometry ended, the ICerPE partner physician and researcher asked him to perform a blood test of the following biomarkers: blood count: [red blood cells (RBCs); hemoglobin (Hb); hematocrit (Hct); mean corpuscular volume (MCV); mean corpuscular hemoglobin (MCH); mean corpuscular hemoglobin concentration (MCHC); total leukocytes; red cell distribution width (RDW)]; serum ferritin (SF); transferrin saturation (TS); vitamin B12; homocysteine; C-reactive protein (CRP); fibrinogen; 25-hydroxy-vitamin-D3 (D3); apolipoprotein A-1 (Apo-A1); apolipoprotein B (Apo-B); glutamic oxalacetic transaminase (GOT); glutamic-pyruvic transaminase (GPT); gamma-glutamyl transpeptidase (Gamma GT); total cholesterol (TC); triglycerides (TG), and; high-density lipoprotein (HDL) and low-density lipoprotein (LDL).

After this stage, the patient went on to confinement due to the COVID-19 pandemic. However, on April 23, 2020, the patient's daughter sent us a message, saying that her father had not respected the confinement recommendations and that, unfortunately, he had died on April 18, 2020.

Results of the assessment by functional neurometry and biomarkers

The result of the patient's anxiety control assessed indirectly by the functional reserve, in percentage, was 7.27% [expected value: 75%] (Table 1). This result is usually compatible with exhaustion of the adrenal gland and/or fatty liver. In a semiological analysis, as the patient was 23.3 kg overweight and with an abdominal circumference of 117 cm, diagnostic compatibility increased in probability. Even though, some thin patients also have hepatic steatosis. The gamma GT level found was 67 [Reference value: 7–45 U/L] (Table 2). The high level of gamma GT, exceeding the laboratory's maximum reference value, is related to insulin resistance, preventing diabetes, and can also signal toxic and/or free radical intoxication, in addition to indicating an affected liver.

The gamma-GT catalyzes a transpeptidation reaction that is involved in the glutathione metabolism. Glutathione is abundant in the epithelial fluid of the lung lining. However, little is known about gamma-GT expression in pulmonary alveoli epithelial cells. There is, however, a relevant study⁴⁹ showing that the pulmonary alveolar epithelial cell type 2 expresses the gene for gamma-GT. The gamma-GT expression in the type 2 pulmonary alveolar epithelial cell is via mRNA III, a transcript that was initially cloned from the liver. This cell synthesizes the gamma-GT protein and releases enzyme activity in a pool associated with surfactant in the pulmonary alveolus.⁴⁹ The results of this study⁴⁹ suggest that the surfactant may play an expanded role in the biology of lung cells as a vehicle for the redistribution of proteins anchored in amphipathic signals across the gas exchange surface of the lung.

The gamma GT transports the glutathione that is outside the cell into the cell and thus keeps the levels of reactive oxygen species low. Some epidemiological studies^{50,51} have shown that gamma GT levels, within laboratory reference values, closest to the highest reference value, are associated with cardiovascular risks, with predictors of future cardiovascular diseases, hypertension, stroke and type II diabetes mellitus.

There is still much to be researched and studied to better understand the relationship between the liver and the lung during deaths caused by COVID-19. However, lung injuries have been considered as the main damage caused by SARS-CoV-2 infection and there are already findings reporting that liver damage occurred during the course of the disease in severe cases.⁵²

Regarding the baroreflex index, the patient presented 60.92% [expected value: >90%] (Table 1) and hemodynamics of 26.07% [expected value: <10%] (Table 1), values found in patients with endothelial dysfunction usually in outpatient consultations. The endothelial dysfunction in the case of the patient in this study seems to have been confirmed by the level of his gamma GT, reported in the above-mentioned paragraphs, of 67 [reference value: 7–45 U/L] (Table 2). There is an epidemiological study in the literature that corroborates this interpretation. This study was carried out with 4,266 participants, demonstrating the association of gamma GT with C-reactive protein in men and women and, mainly, with the stiffening of the brachial artery in men.⁵¹

The endothelial dysfunction in this case report can still be identified by several biomarkers, among which are: low 25-hydroxy-vitamin D3 of 21.6 [reference value: >30] (Table 2), high CRP of 1.74 [reference value: <0.3 mg/dL] (Table 2) and the physiological response of 33.77% [reference value: 31.5–32.5] (Table 2). The digestive overload gives meaning to the low absorption of vitamin B12, found to measure 153 in the patient's blood test [reference value: 210–980 pg/mL] (Table 2). The decrease in vitamin B12 also causes an increase in homocysteine 16.5 [reference value: 5–15 µmol/L] (Table 2), which further increases the probability of endothelial dysfunction.

The patient in this study furthermore had a low 25-hydroxy-vitamin D3 and high C-reactive protein. However, an important finding in the literature⁵³ analyzed the relationship of vitamin D3 and C-Reactive protein with asymptomatic individuals supplemented with 21 ng/mL vitamin D3, and found no significant reduction specifically in C-reactive protein.

However, in different contexts from the previous study, another very relevant study⁵⁴ demonstrated that patients supplemented with vitamin D3 of 10 ng/mL for 3 months showed improvement in endothelial function, reduction of oxidative stress and improvement in insulin sensitivity. This finding may encourage the scientific community to improve respiratory microcirculation, as obstruction and vascular resistance have been a major concern for scientists during the COVID-19 pandemic.⁵⁵

And to further encourage researchers in relation to the benefits of vitamin D, an epidemiological study by Melamed et al. (2008) found that low levels of vitamin D3 increase the risk of mortality. This study used NHANES III data, collected from 1988 to 1994, studying 13,331 adult individuals of both genders with an average follow-up of 8.8 years. The result was 1,806 deaths, 43% from cardiovascular disease, 26% from cancer, 6% from infectious diseases and 5% from external causes. It was found that people with serum vitamin 25-OH-vitamin-D3 levels between 40–49 ng/mL had a lower mortality rate.⁵⁶

The indication of digestive overload found in the patient in this study using the functional neurometry²⁷ may be related to parasites, but it may also have been due to sedentary habits,

Table 1 – Synthesis of the results of the functional neurometry exam of the male patient who died after being affected by COVID-19.

Functional neurometry							
Anxiety control/Functional reserve		Baroreflex index		Hemodynamics		Physiological response	
Expected	Result	Expected minimum	Result	Expected maximum	Result	Expected range	Result
75%	7.27%	90%	60.92%	10%	26.07%	31.5–32.5%	33.77%

Table 2 – Synthesis of the results of the biomarkers of the male patient who died after being affected by COVID-19.

Blood biomarkers							
RBC		Hb		Hct		MCV	
Reference value	Result	Reference value	Result	Reference value	Result	Reference value	Result
4.4–6.6 million/ μ L	4.42	13–18	13.8	39–54%	42.3%	76–96	95.7
MCH		MCHC		Total leukocytes		RDW	
Reference value	Result	Reference value	Result	Reference value	Result	Reference value	Result
27 - 33	31.2	32–36	32.6	4.000–10.000/ μ L	5.430	11–15	13.6
Serum ferritin		Transferrin saturation		B12 vitamin		Homocysteine	
Reference value	Result	Reference value	Result	Reference value	Result	Reference value	Result
23.90 and 336.20 ng/mL	161.2	20–50%	20.8	210.0–980.0 pg/mL	153	5–15 μ mol/L	16.5
C-reactive protein		Fibrinogen		25-Hydroxy-vitamin D3		Apolipoprotein A-1	
Reference value	Result	Reference value	Result	Reference value	Result	Reference value	Result
<0.3 mg/dL	1.74	200–400 mg/dL	395	>30	21.6	104–202 mg/100 mL	142
Apolipoprotein B		Glutamic oxalacetic transaminase		Glutamic-pyruvic transaminase		Gamma GT	
Reference value	Result	Reference value	Result	Reference value	Result	Reference value	Result
66–133 mg/100 mL	113	<34 U/L	17	10–49 U/L	11	7–45 U/L	67
Total cholesterol		Triglycerides		HDL		LDL	
Reference value	Result	Reference value	Result	Reference value	Result	Reference value	Result
<190 mg/dL	200	<150 mg/dL	611	>40 mg/dL	42	<130 – Low Risk	36
CT/HDL ratio		TG/HDL ratio		LDL/HDL ratio		APO B/APO A1 ratio	
Reference value	Result	Reference value	Result	Reference value	Result	Reference value	Result
≤ 3.3	4.76	≤ 2.5	14.4	≤ 2.3	0.85	≤ 0.5	0.7957

which seems to explain his low vitamin B12 as well. This result seems to reinforce the hypotheses about the obstructions of the respiratory microcirculation.⁵⁵

Kwok et al. (2012)⁵⁷ conducted a study using vitamin B12 in 50 vegetarian subjects who had serum B12 <150 pmol/L (<202 pg/mL) and the results showed a significant serum increase in B12, a decrease in homocysteine and a significant improvement in dilation mediated by brachial flow from $6.3 \pm 1.8\%$ to $7.4 \pm 1.7\%$. The conclusion of this study was that B12 supplementation improved the arterial function of vegetarians with subnormal B12 levels, thus proposing a new strategy for the prevention of atherosclerosis.

In this context of importance of vitamin B12, another study⁵⁸ recently published reinforces the importance of vitamin B12 in the treatment of COVID-19, the authors point out that the virus of the Coronaviridae family (COVID-19) has a single-stranded positive RNA genome. This genome encodes the nsp12 protein, which hosts the RNA polymerase-dependent RNA (RdRP) activity responsible for replicating the viral genome.

Thus, the authors⁵⁸ report that a nsp12 homology model was prepared using the SARS nsp12 (6NUR) framework and

that it was used to screen silica to identify molecules among FDA-approved natural products or drugs that could potentially inhibit nsp12 activity. However, this experiment showed that vitamin B12 (methylcobalamin) can bind to the active site as an inhibitor of the nsp12 protein.⁵⁸

Regarding total cholesterol (TC), triglycerides (TG), HDL and LDL, this study highlighted the results of TG and HDL of the patient who died after acquiring COVID-19. In addition to the TG being well above 611 mg/dL (Table 2) of the expected reference value [<150 mg/dL], there are findings in the literature demonstrating ideal values expected from CT/HDL ratios ≤ 3.3 , TG/HDL ≤ 2.5 and LDL/HDL ≤ 2.3 ,⁵⁹ called atherogenicity indexes, functioning as a safe predictor of high lipoprotein density with the risk of coronary disease. Data from the Framingham study show that each 1 mg/dL increase in HDL decreases the coronary disease mortality rate by 2% for men and 3% for women.^{59,60}

However, Tian and Fu (2010)⁵⁹ establish that the greater the result of the relationship between CT/HDL and TG/HDL, the less will be the protective effect of HDL, because it will indicate a greater quantity of small HDLs (HDL Pre 1) and lesser quantities of large HDLs (HDL2b) and add that when these rela-

tionships worsen, not only the HDL amount worsens, but the type of HDL as well.⁵⁹

The HDL is responsible for much more than transporting cholesterol from arteries and peripheral tissues to the liver; it has antioxidant action, anti-inflammatory action, pro-fibrinolytic action, antithrombotic action, and even reverse cholesterol transport.⁶¹ Thus, when the relationships based on the above-mentioned study were replicated in this case report, the following results were found: CT/HDL = 4.76, TG/HDL = 14.4 and LDL/HDL = 0.85, indicating that the patient in this case report had a high level of atherogenicity.

It also seems relevant that, if there is a decision to initiate early care with the population to improve the microvascular state in order to have better conditions to face the COVID-19 virus, it will be necessary to better understand other relationships, such as apolipoprotein A1 (APO-A1) and apolipoprotein B (APO-B).

O'Keefe et al. (2008) measured APO-A1 and saw that APO-A1 does not become a risk factor when in high concentrations.⁶² A large HDL can be an atherogenic HDL because it can donate cholesterol in peripheral tissues through the scavenger receptor class B1 (SR-B1), unless the APO-A1 is also large.⁶²

Thus, it also seems relevant to investigate the relationship between Apo B and Apo A1. In this case report, when the studied patient's APO B [113 mg/100 mL] was divided by his APO A1 [142 mg/100 mL], a result of 0.7957 was found. However, a study by Tian and Fu (2010)⁵⁹ showed that people with an APO B/APO A1 ratio with a value equal to or less than 0.5 die less from cardiovascular disease.

Conclusion

The assessment of the baroreflex and hemodynamic index by functional neurometry by health professionals can anticipate the identification of people who will need to have a blood test to investigate the biomarkers that indicate subclinical inflammation, hepatic steatosis and endothelial dysfunction.

This study found subclinical inflammation and endothelial dysfunction. However, as this study is only a case report, it is suggested to conduct a clinical study with several patients to ensure statistical significance.

Conflicts of interest

The authors declare no conflicts of interest.

REFERENCES

- Gordan R, Gwathmey JK, Xie L-H. Autonomic and endocrine control of cardiovascular function. *World J Cardiol*. 2015;7(4):204.
- Toorop RJ, Ousrouit R, Scheltinga MR, Moll FL, Bleyls RL. Carotid baroreceptors are mainly localized in the medial portions of the proximal internal carotid artery. *Ann Anat Anatomisch Anzeig*. 2013;195(3):248–52.
- Hainsworth R, Ledsome J, Carswell F. Reflex responses from aortic baroreceptors. *Am J Physiol Legat Cont*. 1970;218(2):423–9.
- Kougias P, Weakley SM, Yao Q, Lin PH, Chen C. Arterial baroreceptors in the management of systemic hypertension. *Med Sci Monit: Int Med J Exp Clin Res*. 2010;16(1):RA1.
- Kirchheim HR. Systemic arterial baroreceptor reflexes. *Physiol Rev*. 1976;56(1):100–77.
- McArdle WD, Katch FI, Katch VL. Exercise physiology: nutrition, energy, and human performance. Lippincott: Williams & Wilkins; 2010.
- Thiriet M. Biology and mechanics of blood flows: Part II: Mechanics and medical aspects. Springer Science & Business Media; 2007.
- Houston M. New concepts in the diagnosis and non-surgical treatment of cardiovascular disease. *Intern Med*. 2014;(S):12.
- Heitzer T, Schlinzig T, Krohn K, Meinertz T, Münzel T. Endothelial dysfunction, oxidative stress, and risk of cardiovascular events in patients with coronary artery disease. *Circulation*. 2001;104(22):2673–8.
- Kones R. Primary prevention of coronary heart disease: integration of new data, evolving views, revised goals, and role of rosuvastatin in management. A comprehensive survey. *Drug Des Dev Therapy*. 2011;5:325–80.
- Wildman RP, Muntner P, Chen J, Sutton-Tyrrell K, He J. Relation of inflammation to peripheral arterial disease in the national health and nutrition examination survey, 1999–2002. *Am J Cardiol*. 2005;96(11):1579–83.
- Von Eynatten M, Hamann A, Twardella D, Nawroth PP, Brenner H, Rothenbacher D. Relationship of adiponectin with markers of systemic inflammation, atherogenic dyslipidemia, and heart failure in patients with coronary heart disease. *Clin Chem*. 2006;52(5):853–9.
- Yekutieli M, Brooks M, Ohry A, Yarom J, Carel R. The prevalence of hypertension, ischaemic heart disease and diabetes in traumatic spinal cord injured patients and amputees. *Spinal Cord*. 1989;27(1):58–62.
- Lehto S, Rönnemaa T, Haffner SM, Pyörälä K, Kallio V, Laakso M. Dyslipidemia and hyperglycemia predict coronary heart disease events in middle-aged patients with NIDDM. *Diabetes*. 1997;46(8):1354–9.
- Li G, Chen X, Jang Y, Wang J, Xing X, Yang W, et al. Obesity, coronary heart disease risk factors and diabetes in Chinese: an approach to the criteria of obesity in the Chinese population. *Obesity Rev*. 2002;3(3):167–72.
- Wilhelmsen L. Coronary heart disease: epidemiology of smoking and intervention studies of smoking. *Am Heart J*. 1988;115(1):242–9.
- Abbasi F, Brown BW, Lamendola C, McLaughlin T, Reaven GM. Relationship between obesity, insulin resistance, and coronary heart disease risk. *J Am Coll Cardiol*. 2002;40(5):937–43.
- Rašlová Kn, Smolková B, Vohnout B, Gašparovič J, Frohlich JJ. Risk factors for atherosclerosis in survivors of myocardial infarction and their spouses: comparison to controls without personal and family history of atherosclerosis. *Metabol Clin Exp*. 2001;50(1):24–9.
- Sprague AH, Khalil RA. Inflammatory cytokines in vascular dysfunction and vascular disease. *Biochem Pharmacol*. 2009;78(6):539–52.
- Zernecke A, Weber C. Chemokines in the vascular inflammatory response of atherosclerosis. *Cardiovasc Res*. 2010;86(2):192–201.
- Blankenberg S, Barbaux S, Tiret L. Adhesion molecules and atherosclerosis. *Atherosclerosis*. 2003;170(2):191–203.
- Lindquist S, Craig E. The heat-shock proteins. *Annu Rev Genet*. 1988;22(1):631–77.
- Chan AC, Carter PJ. Therapeutic antibodies for autoimmunity and inflammation. *Nature Rev Immunol*. 2010;10(5):301–16.

24. Salmon JE, Girardi G. Antiphospholipid antibodies and pregnancy loss: a disorder of inflammation. *J Reprod Immunol.* 2008;77(1):51–6.
25. Sies H. What is oxidative stress? Oxidative stress and vascular disease. Springer; 2000. p. 1–8.
26. Motts JA, Shirley DL, Silbergeld EK, Nyland JF. Novel biomarkers of mercury-induced autoimmune dysfunction: a cross-sectional study in Amazonian Brazil. *Environ Res.* 2014;132:12–8.
27. Ribas V, Guerra Ribas R, Viana M, da Silva Dias J, Cavalcanti I, Rodrigues M, et al. The functional neurometry of Nelson Alves Pereira Júnior: an advanced method of mapping and biofeedback training of the autonomic nervous system functions. *J Psychol Psychotherapy Res.* 2020;7:1–19.
28. Braunwald EEA. Harrison's principles of internal medicine. 15th ed. New York: McGraw-Hill; 2001.
29. Nagai Y, Jones CI, Sen A. Galvanic skin response (GSR)/electrodermal/skin conductance biofeedback on epilepsy: a systematic review and meta-analysis. *Front Neurol.* 2019;10:377. PubMed PMID: 31068887. Pubmed Central PMCID: 6491510.
30. Oleson T. Auriculotherapy stimulation for neuro-rehabilitation. *NeuroRehabilitation.* 2002;17(1):49–62. PubMed PMID: 12016347.
31. Schwartz MS, Andrasik F. Biofeedback: a practitioner's guide. Guilford Publications; 2017.
32. Roehrig KL, Allred JB. Direct enzymatic procedure for the determination of liver glycogen. *Anal Biochem.* 1974;58(2):414–21.
33. Shimazu T, Amakawa A. Regulation of glycogen metabolism in liver by the autonomic nervous system. II. Neural control of glycogenolytic enzymes. *Biochim Biophys Acta: Gen Subj.* 1968;165(3):335–48.
34. Hems DA, Whitton PD. Control of hepatic glycogenolysis. *Physiol Rev.* 1980;60(1):1–50.
35. Piorsky AN, Muniz MMF. Resposta neurométrica computadorizada da microfisioterapia: relato de caso. *Rev Bras Neurometr.* 2017;2(1):5–12.
36. VanBavel E, Siersma P, Spaan JA. Elasticity of passive blood vessels: a new concept. *Am J Physiol Heart Circ Physiol.* 2003;285(5):H1986–2000.
37. Levy MN. Brief reviews: sympathetic-parasympathetic interactions in the heart. *Circ Res.* 1971;29(5):437–45.
38. Barcelos DP. Correlação entre o sistema cognitivo e o fisiológico, através da resposta galvânica e a coerência cardíaca na neurometria. *Rev Bras Neurometr.* 2017;1(1):5–14.
39. Madden KM, Lockhart C, Potter TF, Cuff D. Aerobic training restores arterial baroreflex sensitivity in older adults with type 2 diabetes, hypertension, and hypercholesterolemia. *Clin J Sport Med.* 2010;20(4):312.
40. Lantelme P, Khettab F, Custaud M-A, Rial M-O, Joanny C, Gharib C, et al. Spontaneous baroreflex sensitivity: toward an ideal index of cardiovascular risk in hypertension? *J Hyperten.* 2002;20(5):935–44.
41. Carlson JT, Hedner JA, Sellgren J, Elam M, Wallin BG. Depressed baroreflex sensitivity in patients with obstructive sleep apnea. *Am J Respir Crit Care Med.* 1996;154(5):1490–6.
42. Bristow J, Honour A, Pickering T, Sleight P. Cardiovascular and respiratory changes during sleep in normal and hypertensive subjects. *Cardiovasc Res.* 1969;3(4):476–85.
43. Attwell D, Iadecola C. The neural basis of functional brain imaging signals. *Trends Neurosci.* 2002;25(12):621–5.
44. Piccirillo G, Nocco M, Moisé A, Lionetti M, Naso C, di Carlo S, et al. Influence of vitamin C on baroreflex sensitivity in chronic heart failure. *Hypertension.* 2003;41(6):1240–5.
45. Monahan KD, Eskurza I, Seals DR. Ascorbic acid increases cardiovagal baroreflex sensitivity in healthy older men. *Am J Physiol: Heart Circ Physiol.* 2004;286(6):H2113–7.
46. Béchir M, Enseleit F, Chenevard R, Muntwyler J, Lüscher TF, Noll G. Folic acid improves baroreceptor sensitivity in hypertension. *J Cardiovasc Pharmacol.* 2005;45(1):44–8.
47. Ribeiro EE, Martinez Filho EE. Hemodinâmica e cardiologia intervencionista: abordagem clínica; 2008.
48. Crissinger KD, Burney DL. Influence of luminal nutrient composition on hemodynamics and oxygenation in developing intestine. *Am J Physiol: Gastrointest Liver Physiol.* 1992;263(2):G254–60.
49. Joyce-Brady M, Takahashi Y, Oakes SM, Rishi AK, Levine RA, Kinlough CL, et al. Synthesis and release of amphipathic gamma-glutamyl transferase by the pulmonary alveolar type 2 cell. Its redistribution throughout the gas exchange portion of the lung indicates a new role for surfactant. *J Biol Chem.* 1994;269(19):14219–26.
50. Lee D-H, Steffen LM, Jacobs DR Jr. Association between serum γ -glutamyltransferase and dietary factors: the Coronary Artery Risk Development in Young Adults (CARDIA) Study. *Am J Clin Nutr.* 2004;79(4):600–5.
51. Saijo Y, Utsugi M, Yoshioka E, Horikawa N, Sato T, Gong Y, et al. The relationship of gamma-glutamyltransferase to C-reactive protein and arterial stiffness. *Nutr Metabol Cardiovasc Dis.* 2008;18(3):211–9.
52. Xu L, Liu J, Lu M, Yang D, Zheng X. Liver injury during highly pathogenic human coronavirus infections. *Liver Int.* 2020;40(5):998–1004.
53. Amer M, Qayyum R. Relation between serum 25-hydroxyvitamin D and C-reactive protein in asymptomatic adults (from the continuous National Health and Nutrition Examination Survey 2001 to 2006). *Am J Cardiol.* 2012;109(2):226–30.
54. Tarcin O, Yavuz DG, Ozben B, Telli A, Ogunc AV, Yuksel M, et al. Effect of vitamin D deficiency and replacement on endothelial function in asymptomatic subjects. *J Clin Endocrinol metabol.* 2009;94(10):4023–30.
55. Tang N, Bai H, Chen X, Gong J, Li D, Sun Z. Anticoagulant treatment is associated with decreased mortality in severe coronavirus disease 2019 patients with coagulopathy. *J Thromb Haemost.* 2020.
56. Melamed ML, Michos ED, Post W, Astor B. 25-hydroxyvitamin D levels and the risk of mortality in the general population. *Arch Intern Med.* 2008;168(15):1629–37.
57. Kwok T, Chook P, Qiao M, Tam L, Poon Y, Ahuja A, et al. Vitamin B-12 supplementation improves arterial function in vegetarians with subnormal vitamin B-12 status. *J Nutr Health Aging.* 2012;16(6):569–73.
58. Narayanan N, Nair DT. Vitamin B12 may inhibit RNA-dependent-RNA polymerase activity of nsp12 from the SARS-CoV-2 virus; 2020. Preprints.
59. Tian L, Fu M. The relationship between high density lipoprotein subclass profile and plasma lipids concentrations. *Lipids Health Dis.* 2010;9(1):118.
60. Link JJ, Rohatgi A, de Lemos JA. HDL cholesterol: physiology, pathophysiology, and management. *Curr Probl Cardiol.* 2007;32(5):268–314.
61. Davidson MH. Focusing on high-density lipoprotein for coronary heart disease risk reduction. *Cardiol Clin.* 2011;29(1):105–22.
62. O'Keefe JH, Gheewala NM, O'Keefe JO. Dietary strategies for improving post-prandial glucose, lipids, inflammation, and cardiovascular health. *J Am Coll Cardiol.* 2008;51(3):249–55.