

sobrevida das pacientes, consolidando imediato benefício à saúde pública.

Palavras-chave: 18F-FDG PET/CT, Cervical cancer, Diagnóstico, Disseminação, Estadiamento.

<https://doi.org/10.1016/j.htct.2024.04.073>

CONTRIBUTION OF OCCUPATIONAL THERAPY TO THE QUALITY OF LIFE IN PATIENTS WITH COLORECTAL CANCER DURING CHEMOTHERAPY

Carmen Silvia Passos Lima ^a,
Pâmela Coimbra Argenton Puga ^b,
Luciana Campanatti Palhares ^b

^a Divisão de Oncologia Clínica, Departamento de Anestesiologia, Oncologia e Radiologia, Faculdade de Ciências Médicas (FCM), Universidade Estadual de Campinas (UNICAMP), Campinas, SP, Brazil

^b General Hospital, Universidade Estadual de Campinas (Unicamp), Campinas, SP, Brazil

Introduction/Justification: Occupational therapy (OT) has been recommended for hospitalized and outpatient patients, but its effects on occupational roles and quality of life in patients with cancer are uncertain. **Objectives:** This study aimed to evaluate the roles of OT as a therapeutic resource during chemotherapy for patients with colorectal cancer (CRC). **Materials and Methods:** This was a prospective, longitudinal, and quantitative study, carried out in the chemotherapy room of the Clinical Oncology Service of the General Hospital of University of Campinas from February to November 2018. Socio-demographic profile of patients was obtained from medical records, and Occupational Role Identification List, SF-36 Quality of Life Questionnaire and the FACT-F Fatigue Questionnaire were applied to patients before and after the OT intervention by the researcher responsible for the study. Four to six OT sessions were performed, depending on the number of chemotherapy cycles, and each session lasted one hour. The therapeutic interventions were making mandalas, reflecting on songs, playing bingo, playing dominoes and painting boxes. **Results:** Thirty-eight patients with CRC were enrolled in the study and 35 (92%) completed all procedures. The average age of the patients was 59 years, 58% of whom were male, undergoing neoadjuvant, adjuvant, and palliative chemotherapy. After OT intervention, there was a change in the occupational roles reported by patients. There was also a significant improvement in quality of life in the pain and functional capacity domain. Women had changes in social/family well-being and social aspects and the older they were, the lower their functional well-being. **Conclusion:** OT provided a new meaning in the performance of occupational roles and contributed to improving the quality of life of patients with CRC during chemotherapy. **Acknowledgements:** The study was supported by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES).

Keywords: Colorectal cancer, Occupational therapy, Quality of life.

<https://doi.org/10.1016/j.htct.2024.04.074>

PRELIMINARY ANTI-PROLIFERATIVE ACTIVITIES OF A PALLADIUM(II) COMPLEX OVER SQUAMOUS CELL CARCINOMA OF TONGUE

Carmen Silvia Passos Lima ^a,
Tuany Zambroti Candido ^a,
João Ernesto de Carvalho ^b,
Ana Lucia Tasca Gois Ruiz ^b, Pedro Paulo Corbi ^c

^a Divisão de Oncologia Clínica, Departamento de Anestesiologia, Oncologia e Radiologia, Faculdade de Ciências Médicas (FCM), Universidade Estadual de Campinas (UNICAMP), Campinas, SP, Brazil

^b Faculdade de Ciências Farmacêuticas, Universidade Estadual de Campinas (UNICAMP), Campinas, SP, Brazil

^c Instituto de Química, Universidade Estadual de Campinas (UNICAMP), Campinas, SP, Brazil

Introduction/Justification: Oral squamous cell carcinoma is considered one of the most prevalent subtypes of head and neck cancers. Treatments include surgical resection, radiotherapy, and chemotherapy in the cases of patients with advanced squamous cell carcinoma (SCC). Cisplatin, or cis-diamminedichloridoplatinum(II), has been used for treatment of several types of cancer worldwide since 1978 including advanced head and neck SCC. The successful use of cisplatin led to the development of second-generation platinum-based drugs, with emphasis on carboplatin and oxaliplatin, which have been used for cancer treatment worldwide. Nevertheless, patients treated with platinum drugs as cisplatin are subjected to adverse effects as nephrotoxicity, and the search for new chemotherapeutic agents with reduced side effects is crucial. After the discovery of the platinum anticancer drugs, new metal-based compounds of copper, silver, gold, ruthenium, palladium and iridium were synthesized and evaluated as potential anticancer agents. Padeliporfin (Tookad® Soluble) was the first palladium(II) complex used in vascular targeted photochemotherapy for low-risk prostate cancer treatment, which also confirms the potential of use of this metal in the synthesis of new chemotherapeutic agents. In this context, our research group has dedicated efforts in the search of novel gold, silver, platinum and palladium complexes for treatment of cancer, with emphasis on SCC. One of the silver complexes with the anti-inflammatory drug nimesulide recently prepared in our group demonstrated in vitro and in vivo activity over SCC cells. **Objectives:** This study aimed to present the in vitro anti-proliferative activities over SCC of a water-soluble palladium(II) complex containing a cysteine derivative as a chelating ligand. **Materials and Methods:** SCC of tongue (SCC4 and SCC15) and a non-tumoral cell line (HaCat, immortalized keratinocyte) were used in this study. The cells were cultivated following methodology previously described in the literature. **Results:** The palladium(II) complex inhibited proliferation of SCC15 cells with a GI50 (concentration of a drug that reduces cell growth by 50%) of 40.28 µg mL⁻¹ but low selectivity was observed when compared to HaCat cells (GI50: 28.33 µg mL⁻¹). On the other hand, the complex did not inhibit SCC4 cell proliferation (GI50 > 250 µg mL⁻¹). **Conclusion:** The palladium(II) complex seems to be indicated for