

to retrospectively evaluation the patients diagnosed with LCH who were followed up and treated in our clinic. **Methodology:** The data of patients over the age of 18 years who were followed up and treated in Bozyaka Training and Research Hospital Hematology Clinic between 2015-2021 were scanned retrospectively from the hospital system. **Results:** Data of 6 patients were obtained. The mean age of the patients was 33.6. There was no difference between the genders. Pain was the reason for admission in 4 patients and was the most common symptom. While the most frequently involved system was the skeletal system with 5 patients, lung involvement was seen in 2 patients. Vinblastine and prednisolone combination therapy was given to 1 patient, who developed steroid-related avascular necrosis. One patient who was planned for combination treatment **Conclusion:** LCH is a rare disease especially seen in children and young adults. It can involve the skeletal system, lungs, and other organs. The prognosis is often good with excision of the lesion or systemic treatment.

<https://doi.org/10.1016/j.htct.2021.10.1061>

#### PP 52

##### PILOMATRIX CARCINOMA WAS BEYOND ANY EXPECTATIONS: A CASE REPORT OF CARCINOMA CLINICIAN SHOULD BE AWARE OF IT

Abdalraouf Omar<sup>1</sup>, Suad Elnnami<sup>2</sup>,  
Ibtihaj Hassan<sup>3</sup>

<sup>1</sup> Tripoli University Hospital and Albadry policlinic

<sup>2</sup> Tripoli University Hospital and University of Tripoli

<sup>3</sup> Tripoli University Hospital

**Introduction:** Pilomatrix carcinoma is a rare cutaneous tumor derived from follicular matrix with about 150 reported cases and it is considered as locally aggressive tumor with a tendency to recur. In addition to lymph node metastases and poor prognosis, it is non-chemotherapy responding in systemic metastasis. **Method and result:** A 37-year-old Libyan man presented with two large, coalesced nodules in the face measuring about 3\*2 cm and 3\*3 cm, treated as a case of lipoma by surgical excision based on clinical diagnosis, reappearing of larger nodule 9\*6\*4 cm involving almost all the left cheek, surgical excision was done with histopathological features of pilomatrix carcinoma infiltrating to the subcutaneous adipose tissue and deep striated muscle with no clear margins. Conventional radiotherapy by electron beam was started using the linear accelerating machine, with total radiotherapy dose 60 gray (Gy) in 30 fraction for six weeks. No local recurrences, nor lymph node or systemic metastasis since June 2020. **Conclusion:** Pilomatrix carcinoma must always be considered in the differential diagnosis of nodular tumors of the head-and-neck due to high recurrence rate after simple excision. Furthermore, local radiotherapy post incomplete surgical excision is the best adjuvant therapy.

<https://doi.org/10.1016/j.htct.2021.10.1062>

#### PP 53

##### WHAT HAVE WE EXPERIENCED WITH COVID-19 IN PATIENTS WITH HEMATOLOGICAL DISORDERS?

Simge ERDEM<sup>1</sup>, Metban MASTANZADE<sup>1</sup>,  
Mustafa Murat OZBALAK<sup>1</sup>,  
Sevda SERT EKINCI<sup>2</sup>, Dilek Ozden OZLUK<sup>1</sup>,  
Tank Onur TIRYAKI<sup>1</sup>, Beyza OLUK<sup>1</sup>,  
Ipek YONAL HINDILERDEN<sup>1</sup>,  
Mustafa Nuri YENEREL<sup>1</sup>, Meliha NALCACI<sup>1</sup>,  
Tufan TUKEK<sup>2</sup>, Sevgi KALAYOGLU BESISIK<sup>1</sup>

<sup>1</sup> Istanbul University Istanbul Faculty of Medicine,  
Internal Medicine Department/Hematology Division

<sup>2</sup> Istanbul University Istanbul Faculty of Medicine,  
Internal Medicine Department

**Objective:** Patients with cancer are considered highly vulnerable to the COVID-19 disease. However, there are still few data in hematologic patients. Some small studies have shown a high mortality on patients with hematologic malignancies and COVID-19. In this study we aim to report a single center experience of the hematologic patient population with COVID-19 disease. **Methodology:** This single centre, retrospective, cohort study included a total of 111 adult patients (aged ≥18 years) with diagnosis of neoplastic and non-neoplastic hematologic diseases between March 2020 and August 2021. Ethics committee approval was obtained from the Istanbul University Istanbul Faculty of Medicine Clinical Research Ethics Committee. Categorical variables were compared using Pearson's Chi-square test. STATA16-MP was used for the statistical analysis. **Results:** A total of 111 patients (median age:55) with hematologic disease were diagnosed with COVID19. Ninety patients had neoplastic hematologic disorder. Forty-five patients were receiving anti-neoplastic treatment at the time of COVID19 diagnosis. A total of 21 patients (overall mortality rate:19%) died and 19 of them had neoplastic disorder. The malignancy mortality rate was estimated to be 21%. 45 of 90 cases were receiving chemotherapy. Ten of these 45 patients (22%) died due to COVID19 disease. **Conclusion:** In our study the majority of patients who died due to COVID-19 had hematological malignancies. The cytokine storm which affects the clinical outcome in COVID-19 may contribute to dismal prognosis in hematologic malignancies in which cytokine increase is a part of process. Most of the succumbed patients were relapsed refractory multiple line treated which may reflect the immune insufficiency. It seemed COVID-19 progress is mostly poor in hematologic malignancies compared otherwise healthy people.

<https://doi.org/10.1016/j.htct.2021.10.1063>

#### PP 54

##### A RARE CAUSE OF ANEMIA IN ADULTHOOD CONGENITAL DYSERYTHROPETIC ANEMIA

Buğra Sağlam, Murat Albayrak,  
Hacer Berna Afacan Öztürk, Merih Reis Aras,