



Scientific Comment

Hematopoietic stem cell mobilization for autologous transplantation in multiple myeloma patients previously exposed to cyclophosphamide, thalidomide, and dexamethasone: is granulocyte-colony stimulating factor alone enough?☆



Afonso Celso Vigorito*

Universidade Estadual de Campinas (UNICAMP), Campinas, São Paulo, Brazil

The paradigm for multiple myeloma (MM) therapy has evolved markedly in the past decade with the introduction of numerous new drugs and improved patient outcomes.¹

Autologous hematopoietic stem cell transplantation (aHSCT) is widely used as part of first line therapy in the treatment of transplant-eligible patients with MM.² In these patients, hematopoietic stem cell (HSC) mobilization for aHSCT has commonly been performed using cyclophosphamide plus granulocyte-colony stimulating factor (G-CSF) or G-CSF alone.^{3,4} However, the induction regimens should not increase the mobilization failure risk. This concern has been especially pertinent to patients previously exposed to cyclophosphamide or lenalidomide during induction, as these drugs appear to hamper HSC mobilization. In the article that accompanies this comment, Crusoe et al. demonstrate the feasibility of using G-CSF alone to mobilize progenitor cells in MM patients induced with a cyclophosphamide, thalidomide and dexamethasone regimen.⁵ The number of CD34⁺ cells mobilized was assessed after using G-CSF with or without cyclophosphamide.

The retrospective study of Crusoe et al. included eighty-eight MM patients who underwent aHSCT at two Brazilian centers.⁵ Collection of $>2.0 \times 10^6$ CD34⁺ cells/kg was considered sufficient. The group that received cyclophosphamide

collected a higher median number of progenitor cells [3.8 (range: 3.1–4.4) vs. 3.2 (range: 2.3–3.8) – p -value = 0.008]. However, the cyclophosphamide used in mobilization did not show advantages in terms of mobilization, or improved response or survival.

Indeed, we have to consider that cyclophosphamide used for mobilization may have some disadvantages, such as raising the cost of the procedure due to hospitalization, higher toxicity as patients treated with cyclophosphamide require more time for engraftment of platelets and neutrophils, and potentially a higher incidence of post-transplant infections. Eliminating cyclophosphamide from the mobilization regimen has improved patient convenience and has decreased the duration of mobilization treatment by approximately nine days.⁶

We might also argue that the higher number of progenitor cells collected with cyclophosphamide plus G-CSF would enable the storage of more HSC for a second salvage transplant. However, the utilization of stored autologous HSC to support a second aHSCT in MM patients in the era of novel agent therapies has been addressed. Data from Seattle showed that of 726 patients who had residual HSC in storage after their first aHSCT, only 135 patients underwent a second aHSCT.⁷ The percentage of patients receiving a second aHSCT has declined over time. The resources required to collect and store

DOI of original article: <http://dx.doi.org/10.1016/j.bjhh.2016.06.004>.

☆ See paper by Crusoe et al. in Rev Bras Hematol Hemoter. 2016;38(4):302–309.

* Correspondence to: Hemocentro de Campinas, Rua Carlos Chagas, 480, 13083 878 Campinas, SP, Brazil.

E-mail address: afonso@unicamp.br

<http://dx.doi.org/10.1016/j.bjhh.2016.07.006>

1516-8484/© 2016 Associação Brasileira de Hematologia, Hemoterapia e Terapia Celular. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

unused HSC added up to 336 extra patient days of apheresis and 41,587 extra patient months of cryopreservation, translating into a higher average cost per patient. The authors concluded that a reconsideration of conventional HSC collection and storage practices would save significant cost for the majority of MM patients who never undergo a second aHSCT.⁶ We do not know whether the results from Seattle might be translated to the reality of Brazilian patients treated within the Brazilian National Health System (SUS) that, in some centers, precludes the use of novel and more effective agents that would lead to better responses before and after aHSCT.

In conclusion, the study by Crusoe et al. showed that sufficient progenitor cells can be mobilized to perform at least one aHSCT with the use of G-CSF alone in patients induced using the cyclophosphamide, thalidomide and dexamethasone protocol. This strengthens the current study as it makes a valuable judgment regarding the limitations of performing aHSCT in Brazil and the need to cut costs. The questions addressed here will have a practical impact on the clinical practice.

Conflicts of interest

The author declares no conflicts of interest.

REFERENCES

1. Kumar SK, Dispenzieri A, Lacy MQ, Gertz MA, Buadi FK, Pandey S, et al. Continued improvement in survival in multiple myeloma: changes in early mortality and outcomes in older patients. *Leukemia*. 2014;28:1122-8.
2. Shah N, Callander N, Ganguly S, Gul Z, Hamadani M, Costa L, et al. Hematopoietic stem cell transplantation for multiple myeloma: guidelines from the American society for blood and marrow transplantation. *Biol Blood Marrow Transplant*. 2015;21:1155-66.
3. Giralt S, Stadtmauer EA, Harousseau JL, Palumbo A, Bensinger W, Comenzo RL, et al. International myeloma working group (IMWG) consensus statement and guidelines regarding the current status of stem cell collection and high-dose therapy for multiple myeloma and the role of plerixafor (AMD 3100). *Leukemia*. 2009;23:1904-12.
4. Bensinger W, DiPersio JF, McCarty JM. Improving stem cell mobilization strategies: future directions. *Bone Marrow Transplant*. 2009;43:181-95.
5. Crusoe EQ, Higashi F, Martinez GA, Barros JC, Bellesso M, Rossato M, et al. Is it feasible to use granulocyte-colony stimulating factor alone to mobilize progenitor cells in multiple myeloma patients induced with a cyclophosphamide, thalidomide and dexamethasone regimen? *Rev Bras Hematol Hemoter*. 2016;38.
6. Gertz MA, Kumar SK, Lacy MQ, Dispenzieri A, Hayman SR, Buadi FK, et al. Comparison of high-dose CY and growth factor with growth factor alone for mobilization of stem cells for transplantation in patients with multiple myeloma. *Bone Marrow Transplant*. 2009;43:619-25.
7. Phipps C, Linenberger M, Holmberg LA, Green D, Becker P, Connelly-Smith L, et al. Utilization of stored autologous PBSCs to support second autologous transplantation in multiple myeloma patients in the era of novel agent therapy. *Bone Marrow Transplant*. 2015;50:663-7.

1. Kumar SK, Dispenzieri A, Lacy MQ, Gertz MA, Buadi FK, Pandey S, et al. Continued improvement in survival in multiple