

Images in Clinical Hematology

Megakaryocytes in pulmonary circulation: an “old” knowledge with new implications

Leonardo Rodrigues de Oliveira *, Antônio Carlos de Oliveira Meneses 

Universidade Federal do Triângulo Mineiro (UFTM), Uberaba, MG, Brazil



ARTICLE INFO

Article history:

Received 24 December 2020

Accepted 24 February 2021

Available online 24 May 2021

The post-mortem pulmonary findings of a 59-year-old woman with infection by the human immunodeficiency virus and liver cirrhosis related to chronic hepatitis B are presented.




Figure 1 – Multiple megakaryocytes (arrows) identified in the pulmonary circulation (Hematoxylin and Eosin staining, 400 × magnification).

The patient was admitted due to decompensated liver disease and initial laboratory data revealed anemia (hemoglobin concentration 8.8 g/dL), leukocytosis ($12 \times 10^6/L$) and normal platelet count ($399 \times 10^9/L$). The patient was receiving prolonged highly active antiretroviral therapy. No treatment directed at chronic hepatitis B, tuberculosis or anti-cancer therapy has been administered.




Figure 2 – Immunohistochemistry analysis for confirmation of megakaryocytes (arrows) in lungs. Factor VIII-related anti-gen stain. CD61 immunohistochemistry was also positive on megakaryocytes (not shown). 400 × magnification.

* Correspondence author: Rua Getúlio Guaritá, 130, 38025-440 Uberaba, MG, Brazil +55-34-33185046

E-mail addresses: leonardo.oliveira@ebserh.gov.br (L.R. de Oliveira), acomeneses@gmail.com (A.C.d.O. Meneses).
<https://doi.org/10.1016/j.htct.2021.02.012>

2531-1379/© 2021 Published by Elsevier España, S.L.U. on behalf of Associação Brasileira de Hematologia, Hemoterapia e Terapia Celular. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Figure 3 – Bone marrow biopsy shows normal global cellularity (granulocytic hypoplasia and erythroid hyperplasia) and normal megakaryocyte cellularity (2.8 megakaryocytes per high-power field [$\times 400$] – normal range 2 to 4 per high-power field). There is no evidence of cancer infiltration, granulomas, or mycoses (Hematoxylin and Eosin staining, $200\times$ magnification).

In *post-mortem* study, hemoperitoneum was identified as the immediate cause of death in association with active pulmonary tuberculosis and hepatitis B virus-associated hepatocarcinoma (TNM Staging System – IVB). Numerous megakaryocytes were detected in pulmonary circulation

(Figures 1 and 2). In bone marrow evaluation, mycobacterial or fungal infections as well as cancer infiltration were not detected (Figure 3). The evidence of thrombopoiesis in lungs is not recent and innovative experiments indicate that up to half of platelet production may originate from megakaryocytes located in the pulmonary circulation.^{1,2,3}

Conflicts of interest

The authors declare no conflicts of interest.

REFERENCES

1. Levine RF, Eldor A, Shoff PK, Kirwin S, Tenza D, Cramer EM. Circulating megakaryocytes. Delivery of large numbers of intact, mature megakaryocytes to the lungs. *Eur J Haematol.* 1993;51(4):233–46. <https://doi.org/10.1111/j.1600-0609.1993.tb00637.x>.
2. Lefrançois E, Ortiz-Muñoz G, Caudrillier A, Mallavia B, Liu F, Sayah DM, et al. The lung is a site of platelet biogenesis and a reservoir for haematopoietic progenitors. *Nature.* 2017;544(7648):105–9. <https://doi.org/10.1038/nature21706>.
3. Borges I, Sena I, Azevedo P, Andreotti J, Almeida V, Paiva A, et al. Lung as a niche for hematopoietic progenitors. *Stem Cell Rev.* 2017;13(5):567–74. <https://doi.org/10.1007/s12015-017-9747-z>.