

databases using the terms CART therapy, haematological malignancies, and hypogammaglobulinemia. Articles including patients with any haematological malignancies undergone CART therapy and assessment done on hypogammaglobulinemia were included. Following screening and selection of the articles, narrative synthesis, quality assessment, and meta-analysis were conducted **Results:** 1197 citations, 9 were finally included for meta-analysis comprising of 425 patients who were affected due to any haematological malignancies and had undergone CART therapy. The overall incidence rate was 35.35%. In all the studies, hypogammaglobulinemia was managed using IgG. Most of the patients across the studies had infection due to reduction in WBC count. The overall incidence of neutropenia following CART therapy was 59% lymphopenia was 82%, and B-cell aplasia was 49.5%. **Conclusion:** The effective way for management of hypogammaglobulinemia was using IgA antibody. The overall incidence of hypogammaglobulinemia and WBCs was difficult to conclude as majority of the studies were of low and fair quality and were collected at different time points after administration of CART therapy. Thus, good quality clinical trials, open label trials or RCT are required. Hypogammaglobulinemia increases with a decrease in neutrophils, lymphocytes, and B-type cells leading to variable infection.

<https://doi.org/10.1016/j.htct.2022.09.1252>

PP18

RECTAL CANCER DISTANCE TO THE ANAL VERGE AND THE T STAGING: MAGNETIC RESONANCE IMAGING FINDINGS

Abdalraouf Omar¹, Fatma Algledy²,
Najat Amar³, Mufeida Elmsurati⁴

¹ Albadri Polyclinic and Tripoli University Hospital, Tripoli, Libya

² National Center of Disease Control – Libya

³ Oil Clinic, Tripoli, Libya

⁴ Tripoli University Hospital and University of Tripoli, Tripoli, Libya

Objective: This study sought to determine the magnetic resonance imaging (MRI) T staging and the rectal cancer (RC) distance to the anal verge in patients treated in radiotherapy department of Tripoli University Hospital. **Methodology:** An observational study was conducted in Radiotherapy department at Tripoli University Hospital retrospectively from January 1, 2018 to December 31, 2020 for total number of 73 patients whom met the inclusion criteria; 18-year-old or more, male and female with primary RC, T2 or more. distance metastasis or secondary RC were excluded. **Results:** Patients were 38 female and 35 male. Patient less than 50 years old was 25% and 38% was between 50-69 years old. patient at 70 years old or older was 10%. The low rectal cancer, less than 5 cm to the anal verge, is in 38.4% of the patients, with most of the patients at T2 staging (45.5%). While 19.2% was in the mid rectum, 5-10 cm to the anal verge, the T2 was 9%. Regarding the high rectum, more than 10 cm to the anal verge, it was

present in 42.5%, of which 45.5% was in T4b. **Conclusion:** Rectal cancer was less commonly in the mid rectum. in the low rectum it was commonly T2 stage and in high rectum T4b was predominant. Further studies are needed.

<https://doi.org/10.1016/j.htct.2022.09.1253>

PP19

A CASE OF FASCIOLOSIS PRESENTING WITH SEVERE HYPEREOSINOPHILIA

Metban Mastanzade, Arzu Şenol, Alper Koç

Elazığ Fethi Sekin City Hospital

Objective: *Fasciola hepatica* is a parasitic trematode and infects livers of various mammals and rarely infects human liver. Frequently eosinophilia is detected in laboratory findings, but it is generally mild or moderate as with other parasitic infections. Here we present a patient with Fasciolosis as the cause of severe hyper eosiphilia. **Case report:** A 66-year-old female patient presented with weight loss, nausea and abdominal pain for one month. Her physical examination was unremarkable except for mild hepatomegaly. Her laboratory tests were as follows; leukocytes 29900/mm³, eosinophils 21550/mm³ (%71.9), ALP 379 IU/L, LDH 278 IU/L, GGT 53 IU/L, CRP 30 mg/dl. All other etiological tests including primary secondary causes were negative. Abdominal MRI revealed focal patchy nodular lesions. *Fasciola hepatica* IHA (1/2560) was positive. **Results:** After the diagnosis, the patient was administered 2 doses of triclabendazole (10 mg/mg) at 5 day intervals. In the 3rd month of the treatment, the control eosinophil count decreased to 480/mm³, and the patient was free of any symptoms. **Conclusion:** Severe eosinophilia (>5000/mm³) is generally associated with malignant diseases, hyper eosinophilic syndrome or primary hematologic disorders. But it would be useful to consider fasciolosis in hyper eosinophilia patients who are sheep and cattle breeder and present with gastrointestinal system complaints such as jaundice and abdominal pain.

<https://doi.org/10.1016/j.htct.2022.09.1254>

PP 20

RITUXIMAB-INDUCED SEVERE ACUTE THROMBOCYTOPENIA IN A PATIENT WITH SPLENIC MARGINAL ZONE LYMPHOMA

Taha Ulutan Kars, Zahit Furkan Yorgancı,
Osman Yaşkıran, Atakan Tekinalp

Necmettin Erbakan University Meram Faculty of Medicine

Objective: Rituximab, which is widely used in the treatment of B-cell lymphoma, is a chimeric monoclonal antibody directed against the CD20 antigen. Rituximab has many side effects, mainly allergic and neurological. Rituximab may cause thrombocytopenia in the long term after