



Revista Brasileira de Hematologia e Hemoterapia Brazilian Journal of Hematology and Hemotherapy

www.rbhh.org



Letter to the Editor

Zika virus and its implication in transfusion safety



Zika virus (ZIKV), an emerging flavivirus, was initially isolated in 1947 from a Rhesus monkey in Uganda. The first evidence of human transmission of ZIKV was reported in 1952.¹ Only sporadic cases of ZIKV in humans were reported in Africa and Asia before 2007, when significant outbreaks were registered outside these regions – on the islands of Micronesia, French Polynesia and New Caledonia in the Pacific Ocean.² In Brazil, the first autochthonous ZIKV transmission was reported in the northeastern region in May 2015.³ It is believed that ZIKV was introduced in Brazil by asymptomatic travelers during the 2014 World Cup or by the World Sprint Championship canoe race.^{4,5} ZIKV has similar characteristics to Dengue virus in relation to epidemiology and the transmission cycle in urban environments.⁶ Therefore, we should expect that, due to the large distribution of the arthropod vectors (*Aedes aegypti* mosquitoes), the number of outbreaks and symptomatic cases might propagate.

In Brazil, there are 17 officially confirmed ZIKV cases in three states: Bahia, Rio Grande do Norte and São Paulo.⁷ However, with the spread of ZIKV in the country, the World Health Organization (WHO) has already reported ZIKV in 14 Brazilian states including Alagoas, Bahia, Ceará, Maranhão, Mato Grosso, Pará, Paraíba, Paraná, Pernambuco, Piauí, Rio de Janeiro, Rio Grande do Norte, Roraima, and São Paulo.⁸ Currently, Brazil is evidencing a high number of cases of microcephaly, mainly in the northeastern part of the country, which was believed to be linked to ZIKV infection acquired during pregnancy. However, this cannot be confirmed due to insufficient scientific support as ZIKV RNA was found in the amniotic fluid of only two cases. If this condition is found to be related to ZIKV, it will represent a novel clinical manifestation of this emerging virus. Other neurological disorders correlated to ZIKV manifestations include: Guillain-Barré syndrome, encephalitis, myelitis, meningoencephalitis, and optical neuritis.⁹

Since ZIKV is an blood-borne virus transmitted by arthropods, it represents potential risk for transfusion safety. Although the significance of ZIKV to the blood transfusion process and use of blood derivatives is currently unknown, there is a risk that ZIKV could also be transmitted by transfusions. A high number of asymptomatic blood donors was observed during the French Polynesia outbreak. However, several questions remain unclear related to the possible impact

of ZIKV in blood transfusion and transfusion medicine in general:

1. *The potential of ZIKV to cause asymptomatic cases must be elucidated.* Asymptomatic cases of ZIKV among blood donors (up to 74%) have been reported.¹⁰ Therefore, the potential of transmission by transfusion exists. To calculate the transfusion-transmission model, it is important to estimate the incidence of infection and the average duration of viremia.¹¹ However, even at a global level, there is no information about the incidence of infection in the general population and the events related to the viremic phase of ZIKV infection.
2. *Viral load of ZIKV infection.* Viral load during ZIKV infection has previously been measured in blood samples of asymptomatic blood donors (3.40–6.91 log copies/mL).¹² Additionally, viral load has been detected in urine 10 to 20 days after the onset of the disease ($0.7\text{--}220 \times 10^6$ copies/mL).¹³ Therefore, transfusion-transmitted ZIKV infection is possible. However, more detailed studies concerning the quantity and viremia period in asymptomatic individuals are essential.
3. *Effectiveness of transmission via blood transfusion.* Until now, the effectiveness of the transmission of ZIKV via transfusion of blood is unknown. As other flaviviruses, such as Dengue virus 1–4 and West Nile virus,¹¹ can be transmitted by blood transfusion and cause clinical symptoms in blood recipients,¹⁴ the transmission of ZIKV by transfusions seems possible.
4. *Inactivation of ZIKV in blood.* It seems that ZIKV is sensitive to blood pathogen inactivation procedures. The application of amotosalen and ultraviolet A illumination reduces viral titers in plasma and ZIKV does not productively infect cell cultures. In addition, by the second passage in cell cultures, ZIKV RNA becomes undetectable.¹⁵ Therefore, the procedures of pathogen inactivation in blood seem to be effective to inactivate ZIKV and prevent transfusion-transmitted infection.
5. *Molecular diagnosis of ZIKV during screening for blood donation.* In Brazil, no serological or molecular tests for ZIKV have been approved by the Ministry of Health. The diagnosis is currently made by isolation of the virus in cell cultures and molecular detection by in-house polymerase chain

reaction systems in Reference Laboratories. The immediate implantation of molecular tests to screen blood donors at this moment seems unfeasible. This is not only due to the high cost of molecular testing but also to the fact that the pathogenesis of ZIKV infection is not fully understood. Moreover, ZIKV infection is seasonal as more cases are reported during the peak proliferation period of the transmitting arthropod vectors, therefore the implementation of blood donor screening for ZIKV should follow the seasonality of infection.

6. *Conduct and deferral of blood donors from donation.* Brazilian blood banks should intensify the pre-donation screening to minimize the risk of the transmission of ZIKV by transfusions. It is important to pay attention to possible disease risk factors (signs and symptoms of an arboviral disease, travel to endemic areas, etc.) in blood donor candidates. We also recommend that blood banks offer educational programs about ZIKV infection to alert about the potential risk of the transmission of this emerging virus.

Conclusion

ZIKV is an emerging virus in Brazil. In the majority of cases, the infection is asymptomatic however mild to severe clinical symptoms have also been described. The efficiency of the transmission of ZIKV by transfusions is still unknown and additional studies are needed to better evaluate the proportion of asymptomatic blood donors infected by ZIKV, the duration of viremia before clinical signs of acute arboviral infection appear, and the clinical outcomes of ZIKV infection. Until now, the only preventive measures to control ZIKV infection are stringent vector control and individual precautions in respect to mosquito bites.

Conflicts of interest

The authors declare no conflicts of interest.

REFERENCES

1. Dick GW, Kitchen SF, Haddow AJ. Zika virus isolations and serological specificity. *Trans R Soc Trop Med Hyg.* 1952;46(5):509-20.
2. Cao-Lormeau VM, Musso D. Emerging viruses in the Pacific. *Lancet.* 2014;384(9954):1571-2.
3. Zanluca C, de Melo VC, Mosimann AL, Dos Santos GI, Dos Santos CN, Luz K. First report of autochthonous transmission of Zika virus in Brazil. *Mem Inst Oswaldo Cruz.* 2015;110(4):569-72.
4. Salvador FS, Fujita DM. Entry routes for Zika virus in Brazil after 2014 world cup: new possibilities. *Travel Med Infect Dis.* 2016;14(1):49-51.
5. Musso D. Zika virus transmission from French Polynesia to Brazil. *Emerg Infect. Dis.* 2015;21(10):1887.
6. Musso D, Cao-Lormeau VM, Gubler DJ. Zika virus: following the path of dengue and chikungunya. *Lancet.* 2015;386(9990):243-4.
7. Rodriguez-Morales AJ. Zika: the new arbovirus threat for Latin America. *J Infect Dev Ctries.* 2015;9(6):684-5.
8. World Health Organization. Epidemiological Update Zika virus infection [Internet]; 2015. Available from: http://www.paho.org/hq/index.php?option=com_docman&task=doc.view&Itemid=270&gid=32021&lang=en [cited 18.12.15].
9. Mallet HP. Bilan de l'épidémie à virus zika en Polynésie française, 2013-2014. *Bulletin d'information sanitaires, épidémiologiques et statistiques* [Internet], vol. 13; 2015. p. 1-5. Available from: http://www.hygiene-publique.gov.pf/IMG/pdf/no13_-mai.2015_-zika.pdf [cited 18.12.15].
10. Musso D, Nhan T, Robin E, Roche C, Bierlaire D, Zisou K, et al. Potential for Zika virus transmission through blood transfusion demonstrated during an outbreak in French Polynesia, November 2013 to February 2014. *Euro Surveill.* 2014;19(14), pii:20761.
11. Petersen LR, Busch MP. Transfusion-transmitted arboviruses. *Vox Sang.* 2010;98(4):495-503.
12. Lanciotti RS, Kosoy OL, Laven JJ, Velez JO, Lambert AJ, Johnson AJ, et al. Genetic and serologic properties of Zika virus associated with an epidemic, Yap State, Micronesia, 2007. *Emerg Infect Dis.* 2008;14(8):1232-9.
13. Gourinat AC, O'Connor O, Calvez E, Goarant C, Dupont-Rouzeyrol M. Detection of Zika virus in urine. *Emerg Infect Dis.* 2015;21(1):84-6.
14. Levi JE, Nishiya A, Félix AC, Salles NA, Sampaio LR, Hangai F, et al. Real-time symptomatic case of transfusion-transmitted dengue. *Transfusion.* 2015;55(5):961-4.
15. Aubry M, Richard V, Green J, Brout J, Musso D. Inactivation of Zika virus in plasma with amotosalen and ultraviolet A illumination. *Transfusion.* 2016;56(1):33-40.

Simone Kashima*, Svetoslav Naney Slavov,
Dimas Tadeu Covas

Universidade de São Paulo (USP), Ribeirão Preto, SP, Brazil

*Corresponding author at: Laboratório de Biologia Molecular, Fundação Hemocentro de Ribeirão Preto, Rua Tenente Catão Roxo, 2501, 14051-140 Ribeirão Preto, SP, Brazil.

E-mail address: skashima@hemocentro.fmrp.usp.br (S. Kashima).

Received 18 December 2015

Accepted 6 January 2016

Available online 4 February 2016

<http://dx.doi.org/10.1016/j.bjhh.2016.01.002>

1516-8484/© 2016 Associação Brasileira de Hematologia, Hemoterapia e Terapia Celular. Published by Elsevier Editora Ltda. All rights reserved.